

# PH-107: Introduction to Quantum Mechanics

## Tutorial Sheet 10

\* marked problems will be solved in the tutorial class (D3-D4: Wednesday, D1-D2: Saturday)

### Simple Harmonic Oscillator and 2D/3D Systems:

1. Using the uncertainty principle, show that the lowest energy of an oscillator is  $\hbar\omega/2$ .
2. Determine the expectation value of the potential energy for a quantum harmonic oscillator (with mass  $m$  and frequency  $\omega$ ) in the ground state. Use this to calculate the expectation value of the kinetic energy. The ground state wavefunction of quantum harmonic oscillator is:

$$\psi_0(x) = C_0 \exp\left(-\frac{m\omega}{2\hbar}x^2\right) \quad C_0 \text{ is constant;} \quad (1)$$

3. A diatomic molecule behaves like a quantum harmonic oscillator with the force constant  $k = 12Nm^{-1}$  and mass  $m = 5.6 * 10^{-26}kg$ 
  - (a) What is the wavelength of the emitted photon when the molecule makes the transition from the third excited state to the second excited state ?
  - (b) Find the ground state energy of vibrations for this diatomic molecule.
4. Vibrations of the hydrogen molecule can be modeled as a simple harmonic oscillator with the spring constant  $k = 1.13 * 10^3 Nm^{-2}$  and mass  $m = 1.67 * 10^{27} kg$ .
  - (a) What is the vibrational frequency of this molecule ?
  - (b) What are the energy and the wavelength of the emitted photon when the molecule makes transition between its third and second excited states ?
5. \* A two-dimensional isotropic harmonic oscillator has the Hamiltonian

$$H = -\frac{\hbar^2}{2m} \left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2}k(x^2 + y^2)$$

- (a) Show that the energy levels are given by

$$E_{n_x, n_y} = \hbar\omega(n_x + n_y + 1) \quad \text{where} \quad n_x, n_y \in (0, 1, 2, \dots) \quad \omega = \sqrt{\frac{k}{m}}$$

- (b) What is the degeneracy of each level?
6. \* Consider the Hamiltonian of a two-dimensional anisotropic harmonic oscillator ( $\omega_1 \neq \omega_2$ )

$$H = \frac{p_1^2}{2m} + \frac{p_2^2}{2m} + \frac{1}{2}m\omega_1^2 q_1^2 + \frac{1}{2}m\omega_2^2 q_2^2$$

- (a) Exploit the fact that the Schrödinger eigenvalue equation can be solved by separating the variables and find a complete set of eigenfunctions of  $H$  and the corresponding eigenvalues.
- (b) Assume that  $\frac{\omega_1}{\omega_2} = \frac{3}{4}$ . Find the first two degenerate energy levels. What can one say about the degeneracy of energy levels when the ratio between  $\omega_1$  and  $\omega_2$  is not a rational number.
7. A particle of mass  $m$  is confined to move in the potential  $(m\omega^2 x^2)/2$ . Its normalized wave function is

$$\psi(x) = \left(\frac{2\beta}{\sqrt{3}}\right) \left(\frac{\beta}{\pi}\right)^{1/4} x^2 e^{-(\beta x^2/2)}$$

where  $\beta$  is a constant of appropriate dimension.

- (a) Obtain a dimensional expression for  $\beta$  in terms of  $m, \omega$  and  $\hbar$ .
- (b) It can be shown that the above wave function is the linear combination

$$\psi(x) = a\psi_0(x) + b\psi_2(x)$$

where  $\psi_0(x)$  is the normalized ground state wave function and  $\psi_2(x)$  is the normalized second excited state wave function of the potential. Evaluate  $b$  and hence calculate the expectation value of the energy of the particle in this state  $\psi(x)$ .

Given:  $I_0(\beta) = \int_{-\infty}^{+\infty} e^{-\beta x^2} dx = \sqrt{\frac{\pi}{\beta}}$ ,  $I_n(\beta) = \int_{-\infty}^{+\infty} (x^2)^n e^{-\beta x^2} dx = (-1)^n \frac{\partial^n}{\partial \beta^n} (I_0(\beta))$ ,

$$\psi_0(x) = \left(\frac{\beta}{\pi}\right)^{1/4} e^{-\frac{\beta x^2}{2}}$$

8. Consider an 3D isotropic harmonic oscillator show that the degeneracy  $g_n$  of the  $n$ th excited state, which is equal the number of ways the non negative integers  $n_x, n_y, n_z$  may be chosen to total to  $n$ , is given by

$$g_n = \frac{1}{2}(n+1)(n+2)$$

9. \* A charged particle of mass '  $m$  ' and charge '  $q$  ' is bound in a 1-dimensional simple harmonic oscillator potential of angular frequency '  $\omega$  '. An electric field  $E_0$  is turned on.
- (a) What is the total potential  $V(x)$  experienced by the charge ?
- (b) Express the total potential in the form of an effective harmonic oscillator potential.
- (c) Sketch  $V(x)$  versus  $x$ .
- (d) What is the ground state energy of the particle in this potential?
- (e) What is the expectation value of the position ( $x$ ) if the charge is in its ground state ?